



Journal of Endovascular Therapy – The use of stents has improved management and outcomes of coronary artery disease, and clinical trials are attempting to prove the same will be true for superficial femoral artery disease. Randomized trials have shown favorable results for self-expanding nitinol stents compared with balloon angioplasty. A

[new report seeks to test this treatment](#)

in a real-world population of patients enrolled in an observational registry.

The [Journal of Endovascular Therapy](#) offers a prospective multicenter observational study of patients with symptomatic superficial femoral artery stenosis. A total of 998 patients suffering from chronic limb ischemia were enrolled at 13 German medical centers. Twelve-month outcomes of the participants were assessed.

The median lesion length was 8 cm, but there was no restriction on the length of lesion to be treated. The authors cite this criterion as supporting the real-world aspect of the study population. Rather than including primarily high-risk patients, a general mix of patients participated in the study. However, a third of the patients had diabetes, and 43 percent were current smokers.

In all, 1,050 lesions were treated. Twelve months after surgery, target lesion revascularization—the primary endpoint of the study—was achieved in 136, or 17 percent, of the patients. Restenosis occurred in 24 percent of the patients and reocclusion in 10 percent. These results indicate that stent use is safe and is delivering favorable outcomes at the 1-year mark.

A [commentary](#) in the same issue asserts that the “real-world” study group is actually representative of a narrower set of patients, largely because the median lesion length was less than 10 cm—a length that has previously shown good success with intervention. The author notes that, although the study validates that superficial femoral artery stenting is producing satisfactory patency results, patient quality of life and functionality measures might better represent success.

Full text of the article, “[German Multicenter Real-World Registry of Stenting for Superficial](#)

Femoral Artery Disease: Clinical Results and Predictive Factors for Revascularization

" and

commentary article

,

Journal of Endovascular Therapy

, Vol. 21, No. 4, 2014, are now available.

###

About the *Journal of Endovascular Therapy*

The *Journal of Endovascular Therapy*, an official publication of the International Society of Endovascular Specialists, publishes peer-reviewed articles of interest to clinicians and researchers in the field of endovascular interventions. The scope of the journal is multidisciplinary, representing all topics related to minimally invasive peripheral vascular diagnosis and treatment. Original clinical studies, experimental investigations, state-of-the-art reviews, rapid communications, case reports, technical notes, editorials, and letters to the editor are published, as well as feature articles on the basics of endovascular interventions.

The journal is available online at

www.jevtonline.org .

To learn more about the society, please visit

www.isesonline.org

.