



Okayama, 5 October - Okayama University researchers control the timing and location of protein synthesis using a photoresponsive compound that is an inactive key molecule until it is activated by brief irradiation.

The production of proteins at distinct times and locations regulates cell functions, such as cell development and differentiation into specific cell types. The ability to exert spatiotemporal control over protein synthesis would help investigate these processes. Now Takashi Ohtsuki and colleagues in the Department of Medical Bioengineering at Okayama University, Japan, have shown that they can prevent protein synthesis reactions from taking place using a photoresponsive molecular cage (see Figure). Brief exposure to light releases key protein synthesis molecules from the cage without damaging them, so that protein synthesis takes place at the time and location of irradiation.

In their study, the researchers used an aminoacyl-transfer RNA (aa-tRNA), a molecule that helps decode messenger RNA so that protein synthesis can occur. Previous work had successfully inhibited protein synthesis by protecting aa-tRNA with a molecular cage of nitroveratryloxycarbonyl group. However the 30 minutes of light exposure required to release the aa-tRNA from the cage caused damage that then inhibited the protein's function.

Instead Ohtsuki's team used (7-diethylaminocoumarin-4-yl)methoxycarbonyl (DEACM) group. While the caged aa-tRNA was stable for at least 4 hours, irradiating for just 20 seconds released the aa-tRNA. The researchers demonstrated light-controlled activity of the caged aa-tRNA *in vitro*, in a gel, in liposomes and in mammalian cells.

"This method of spatiotemporally photocontrolling translation offers a promising approach for investigating the relationship between local translation and biological functions," conclude the researchers in their report of the work. As well as a tool for protein synthesis investigations the team suggest it could also be used to insert artificial amino acids into proteins in a controlled manner for further studies.

Background

What are proteins?

Proteins are biomolecules comprising specific sequences of amino acids. They are responsible for a number of functions including metabolic reaction catalysis, DNA replication, responses to stimuli and molecular transport. Constant protein generation is required to balance the loss of proteins through degradation and secretion from cells.

The synthesis of proteins in specific locations is important in for instance the development of the fly *Drosophila*, where localisation of *oskar* mRNA in the egg determines the anterior-posterior axis. Neuron function also relies on the synthesis of proteins in specific locations.

How are proteins synthesised?

The usual mode for protein synthesis uses transfer RNA (tRNA), messenger RNA (mRNA) and ribosomes to assemble the amino acid chains in the right order. The role of the tRNA is to link the amino acids to the mRNA to help decode it. The ribosome matches the aminoacyl-tRNA (tRNA carrying an amino acid) with the mRNA so that the protein encoded on the mRNA is synthesized.

How does the light-activated DEACM cage work?

The DEACM group protects the aa-tRNA so that it cannot interact with the molecules necessary for protein synthesis. On exposure to light, the DEACM-aa-tRNA degrades into separate components releasing the aa-tRNA.

The researchers demonstrated the light-responsive aa-tRNA release by irradiating a gel containing green fluorescent protein mRNA and the complex through a mask in the shape of a smiley face. No fluorescence was observed until the gel was irradiated with 405nm (blue) light, soon after which fluorescence was observed simulating the shape of the mask.

As well as showing light responsive protein synthesis *in vitro*, gels and liposomes, the researchers also inserted the caged aa-tRNA in Chinese hamster ovary cells. Following 5 hours of incubation, the researchers observed fluorescence of photo-dependently synthesized DsRed protein in the irradiated cells (see following Figure).

Caption

Photoinduced DsRed synthesis in a mammalian cell. DsRed was synthesized only within the irradiated cell.

Reference

Takashi Ohtsuki, Shigeto Kanzaki, Sae Nishimura, Yoshio Kunihiro, Masahiko Sisido, Kazunori Watanabe. Phototriggered protein syntheses by using (7-diethylaminocoumarin-4-yl) methoxycarbonyl-caged aminoacyl tRNAs. *Nature Communications* 7, Article number: 12501 (2016)

DOI : 10.1038/ncomms12501

<http://www.nature.com/ncomms/2016/160817/ncomms12501/full/ncomms12501.html>

Correspondence to

Professor Takashi Ohtsuki, Ph.D.

Department of Medical Bioengineering,

Okayama University, 3-1-1 Tsushimanaka, Kita-ku,

Okayama 700-8530, Japan

e-mail : ohtsuk@okayama-u.ac.jp

Further information

Okayama University

1-1-1 Tsushima-naka , Kita-ku , Okayama 700-8530, Japan

Public Relations and Information Strategy

E-mail: www-adm@adm.okayama-u.ac.jp

Website: http://www.okayama-u.ac.jp/index_e.html

Okayama Univ. e-Bulletin: <http://www.okayama-u.ac.jp/user/kouhou/ebulletin/>

Okayama Univ. e-Bulletin (PDF Issues): <http://www.okayama-u.ac.jp/en/tp/cooperation/ebulletin.html>

About Okayama University (You Tube):

<https://www.youtube.com/watch?v=iDL1coqPRYI>

Okayama University Image Movie (You Tube):

https://www.youtube.com/watch?v=_WnbJVk2eIA

<https://www.youtube.com/watch?v=KU3hOIXS5kk>

Okayama University Medical Research Updates □ OU-MRU □

Vol.1 □ [Innovative non-invasive 'liquid biopsy' method to capture circulating tumor cells from blood samples for genetic testing](#)

Vol.2 □ [Ensuring a cool recovery from cardiac arrest](#)

Vol.3 [Organ regeneration research leaps forward](#)

Vol.4 [Cardiac mechanosensitive integrator](#)

Vol.5 [Cell injections get to the heart of congenital defects](#)

Vol.6 [Fourth key molecule identified in bone development](#)

Vol.7 [Anticancer virus solution provides an alternative to surgery](#)

Vol.8 [Light-responsive dye stimulates sight in genetically blind patients](#)

Vol.9 [Diabetes drug helps towards immunity against cancer](#)

Vol.10 [Enzyme-inhibitors treat drug-resistant epilepsy](#)

Vol.11 [Compound-protein combination shows promise for arthritis treatment](#)

Vol.12 [Molecular features of the circadian clock system in fruit flies](#)

Vol.13 [Peptide directs artificial tissue growth](#)

Vol.14 [Simplified boron compound may treat brain tumours](#)

Vol.15 [Metamaterial absorbers for infrared inspection technologies](#)

Vol.16 [Epigenetics research traces how crickets restore lost limbs](#)

Vol.17 [Cell research shows pathway for suppressing hepatitis B virus](#)

Vol.18 [Therapeutic protein targets liver disease](#)

Vol.19 [Study links signalling protein to osteoarthritis](#)

Vol.20 [Lack of enzyme promotes fatty liver disease in thin patients](#)

Vol.21 [Combined gene transduction and light therapy targets gastric cancer](#)

Vol.22 [Medical supportive device for hemodialysis catheter puncture](#)

Vol.23 [Development of low cost oral inactivated vaccines for dysentery](#)

Vol.24 [Sticky molecules to tackle obesity and diabetes](#)

Vol.25 [Self-administered aroma foot massage may reduce symptoms of anxiety](#)

Vol.26 [Protein for preventing heart failure](#)

Vol.27 [Keeping cells in shape to fight sepsis](#)

Vol.28 [Viral-based therapy for bone cancer](#)

About Okayama University

Okayama University is one of the largest

comprehensive universities in Japan with

roots going back to the Medical Training

Place sponsored by the Lord of Okayama

and established in 1870. Now with 1,300

faculty and 14,000 students, the University

offers courses in specialties ranging from

medicine and pharmacy to humanities

and physical sciences.

Okayama University is located in the heart of Japan approximately 3 hours west of Tokyo by Shinkansen